arXiv:1505.04816v1 [math.AT] 18 May 2015

RATIONAL MODELS OF THE COMPLEMENT OF A SUBPOLYHEDRON IN A MANIFOLD WITH BOUNDARY

HECTOR CORDOVA BULENS, PASCAL LAMBRECHTS,
AND DON STANLEY

Abstract. Let W be a compact simply connected triangulated manifold with boundary and $K \subset W$ be a subpolyhedron. We construct an algebraic model of the rational homotopy type of $W\backslash K$ out of a model of the map of pairs $(K, K\cap\partial W) \hookrightarrow (W, \partial W)$ under some high codimension hypothesis.

We deduce the rational homotopy invariance of the configuration space of two points in a compact manifold with boundary under 2-connectedness hypotheses. Also, we exhibit nice explicits models of these configuration spaces for a large class of compact manifolds.

1. Introduction

Let W be a compact and simply-connected manifold with boundary (in this paper all manifolds are triangulated). Let $f\colon K \hookrightarrow W$ be the inclusion of a subpolyhedron. The first goal of this paper is to determine the rational homotopy type of the complement $W\backslash K$. We will then apply this to deduce the rational homotopy type of the configuration space of two points in a manifold with boundary under 2-connectedness hypotheses. Hence this paper extends the results of [9] and [8] to the case of manifolds *with boundary*.

The main result of [9] is an explicit description of the rational homotopy type of $W\backslash K$ when W is a *closed* manifold and K is a subpolyhedron of codimension $\geq (\dim W)/2+2$. This rational homotopy type depends only on the rational homotopy class of the inclusion $K \hookrightarrow W$ ([9, Theorem 1.2]).

The situation for manifolds with boundary is different. For example, let W be an n-dimensional disk D^n and K be a point. If K is embedded in the interior of D^n then $W\backslash K \simeq S^{n-1}$. On the contrary, if K is embedded in the boundary of D^n then $W\backslash K \simeq *$. Hence the complements $W\backslash K$ have different rational homotopy types, although the two inclusions $K \hookrightarrow W$ are homotopic. These examples show that

Date: 18 May 2015.

2010 *Mathematics Subject Classification.* 55P62, 55R80.

Key words and phrases. Lefschetz duality, Sullivan model, Configuration spaces.

H.C.B. and P.L. are supported by the belgian federal fund PAI Dygest.

we need more information to determine the rational homotopy type of $W\backslash K$. Our main result is that the only extra information needed is related to the inclusion of $\partial W \cap K$ in ∂W. More precisely, we have the following result

Theorem 1.1 (Corollary 4.6 and Theorem 4.5)**.** *Let W be a compact simply connected triangulated manifold with boundary and let K be a subpolyhedron in W.*

Assume that

$$\dim W \geq 2 \dim K + 3. \tag{1}$$

Then the rational homotopy type of $W\backslash K$ depends only on the rational homotopy type of the square of inclusions

$$\begin{array}{ccc} (K \cap \partial W) & \hookrightarrow & \partial W \\ \downarrow & & \downarrow \\ K & \hookrightarrow & W. \end{array} \tag{2}$$

Moreover a CDGA model of $W\backslash K$ (that is, an algebraic model in the sense of Sullivan of this rational homotopy type, see Section 2.1) can be explicitely constructed out of any CDGA model of Diagram (2)*.*

Actually we will see that the high codimension hypothesis (1) can be weakened. Indeed we will establish a sharp *unknotting condition*, which is an inequality relating the connectivity of the inclusion maps and the dimensions of the manifold and the subpolyhedron (see (22) in Corollary 4.6), under which we still get a CDGA model of the complement.

There is an interesting application of this theorem to the study of configuration spaces of 2 points in W,

$$\mathrm{Conf}(W, 2) \coloneqq \{(x_1, x_2) \in W \times W \colon x_1 \neq x_2\}.$$

Indeed this configuration space is the complement

$$\mathrm{Conf}(W, 2) = W \times W \setminus \Delta(W)$$

where $\Delta \colon W \hookrightarrow W \times W$ is the diagonal embedding. We will deduce from Theorem 1.1 the following result.

Theorem 1.2 (Corollary 5.5)**.** *Let W be a 2-connected compact manifold with a 2-connected or empty boundary. The rational homotopy type of the configuration space* $\mathrm{Conf}(W, 2)$ *depends only on the rational homotopy type of the pair $(W, \partial W)$.*

In [4] we prove that a large class of compact manifolds with boundary admit CDGA models of a special form that we call *surjective pretty models*. This class contains in particular even-dimensional disk bundles over a closed manifold and complements of high codimensional polyhedra in closed manifolds. As a consequence, such manifolds admit a

CDGA model of the form P/I where P is a Poincaré duality CDGA and I is some differential ideal. Poincaré duality CDGAs come with a natural *diagonal class* $\Delta \in (P \otimes P)^n$. We then get the following elegant model for the configuration space (see Section 5.3 for more details)

Theorem 1.3. *(Theorem 5.8) Let W be a compact manifold of dimension n with boundary and assume that W and ∂W are 2-connected. If $(W, \partial W)$ admits a surjective pretty model in the sense of* [4] *then a CDGA model of* $\mathrm{Conf}(W, 2)$ *is given by*

$$(P/I \otimes P/I) \oplus_{\overline{\Delta^!}} ss^{-n} P/I,$$

where P is the Poincaré duality CDGA and I the ideal associated to the pretty model, and $\overline{\Delta^!}$ is a map induced by multiplication by the diagonal class $\Delta \in (P \otimes P)^n$.

When W is a closed manifold, we have $I = 0$ and the model of Theorem 1.3 is exactly that of [8].

In the paper [3] in preparation we will show how to build a model (of dgmodules) of $\mathrm{Conf}(W, k)$, $k \geq 2$, which enables to compute effectively the homology of the space of configurations of any number of points in a manifold with boundary. This model will be of the form

$$\left(\frac{(P/I)^{\otimes k} \otimes \Lambda(g_{ij} : 1 \leq i < j \leq k)}{(\text{Arnold and symmetry relations})} \, , \, d(g_{ij}) = \pi_{ij}^*(\overline{\Delta}) \right),$$

mimicking the model in [11].

Here is the plan of the paper. Section 2 contains a very short review on rational homotopy theory, the notion of truncation of a CDGA, a discussion on CDGA structures on mapping cones, and the notion of homotopy kernel. Section 3 is a first step to the understanding of a dgmodule model of the complement $W \setminus K$ and in Section 4 we establish a CDGA model of that complement. In Section 5 we apply the previous results to the model of the configuration space of 2 points in compact manifolds, with some developments of the examples of configuration spaces on a disk bundle or in the complement of a polyhedron in a closed manifold.

2. Truncation of dgmodules and CDGA's, and CDGA structures on mapping cones.

This section contains a quick review on some classical topics that we will need with some special development. In particular in 2.3 we explain some notion of truncation of a CDGA, and in 2.4 we show how to endow a mapping cone (or its truncation) with the structure of a CDGA.

2.1. **Rational homotopy theory.** In this paper we will use the standard tools and results of rational homotopy theory, following [6]. Recall that A_{PL} is the Sullivan-de Rham functor and that for a 1-connected space of finite type, X, $A_{PL}(X)$ is a commutative differential graded algebra (CDGA for short), which completely encodes the rational homotopy type of X. Any CDGA weakly equivalent to $A_{PL}(X)$ is called a *CDGA model of* X. All our dgmodules and CDGAs are over the field $\mathbb{Q}$.

2.2. **Truncation of a dgmodule.** The classical truncation of a cochain complex, i.e. $\mathbb{Q}$-dgmodule, C, is classicaly defined by (see [13, Section 1.2.7])

$$(\hat{\tau}^{\leq N}C)^i = \begin{cases} C^i & \text{if } i < N \\ C^N \cap \ker d & \text{if } i = N \\ 0 & \text{if } i > N \end{cases} \tag{3}$$

This comes with an inclusion

$$\iota\colon \hat{\tau}^{\leq N}C \hookrightarrow C$$

which induces isomorphisms $H^i(\iota)$, for $i \leq N$, and such that $H^{>N}(\hat{\tau}^{\leq N}C) = 0$.

When R is an A-dgmodule, the truncation $\hat{\tau}^{\leq N}R$ is not necessarily an A-dgmodule. In that case a better replacement would be to take for the truncation a quotient R/I where I is a suitable A-dgsubmodule such that $I^i = R^i$ for $i > N$. In this paper we will use the following:

Definition 2.1. *Let R be an A-dgmodule and let N be a positive integer. A* truncation below degree N of R *is an A-dgmodule, $\tau^{\leq N}R$, and a morphism $\pi\colon R \to \tau^{\leq N}R$ of A-dgmodules verifying the two following conditions:*

(1) $(\tau^{\leq N}R)^{>N} = 0$ and $(\tau^{\leq N}R)^{<N} \cong R^{<N}$, and
(2) the morphism π is a surjection of A-dgmodules such that $H^i(\pi)$ is an isomorphism for $0 \leq i \leq N$.

Contrary to $\hat{\tau}^{\leq N}$ from (3), our truncation $\tau^{\leq N}R$ is not unique and is not a functorial construction.

2.3. **Truncation of a CDGA.**

Definition 2.2. *Let A be a connected CDGA.* A CDGA truncation below degree N of A *is a truncation of A-dgmodule $(\tau^{\leq N}A, \pi)$ such that $\tau^{\leq N}A$ is a CDGA and $\pi : A \to \hat{\tau}^{\leq N}A$ is a CDGA morphism.*


Equivalently a CDGA truncation can be seen as a projection $\pi\colon A \to A/I$ where I is an ideal of A such that $I^{<N} = 0$, $I^{>N} = A^{>N}$ and $I^N \oplus (\ker d \cap A^N) = A^N$.

Proposition 2.3. *Any two CDGA truncations below degree N of a given connected CDGA are weakly equivalent.*

Proof. Let A be a connected CDGA and $N \in \mathbb{N}$. It is easy to construct a relative Sullivan model

$$\iota\colon A \rightarrowtail (A \otimes \Lambda V, D)$$

such that $H^{\leq N}(\iota)$ is an isomorphism, $V = V^{\geq N}$ and $H^{>N}(A \otimes \Lambda V, D) = 0$. Indeed, one builds inductively $V = V^{\geq N}$ by adding generators to eliminate all the homology in degrees $> N$. It is straightforward to check that any CDGA truncation $\pi\colon A \to \tau^{\leq N} A$ factors as follows



where $m(V) = 0$. Since $H^{\leq N}(\iota)$ and $H^{\leq N}(\pi)$ are isomorphisms and $H^{>N}(A \otimes \Lambda V, D) = H^{>N}(\tau^{\leq N} A) = 0$, we deduce that m is a quasi-isomorphism. Therefore any two truncation of A are quasi-isomorphic to $(A \otimes \Lambda V, D)$, and hence are weakly equivalent

□

2.4. **Semi-trivial C(D)GA structures on mapping cones.** Let A be a CDGA and let R be an A-dgmodule. We will denote by $s^k R$ the k-th suspension of R, i.e. $(s^k R)^p = R^{k+p}$, and for a map of A-dgmodules, $f\colon R \to Q$, we denote by $s^k f$ the k-th suspension of f. Furthermore, we will use $\#$ to denote the linear dual of a vector space, $\#V = hom(V, \mathbb{Q})$, and $\#f$ to denote the linear dual of a map f.

If $f\colon Q \to R$ is an A-dgmodule morphism, *the mapping cone* of f is the A-dgmodule

$$C(f) \coloneqq (R \oplus_f sQ, \delta)$$

defined by $R \oplus sQ$ as an A-module and with a differential δ such that $\delta(r, sq) = (d_R(r) + f(q), -sd_Q(q))$.

When $R = A$, the mapping cone $C(f\colon Q \to A)$ can be equipped with a unique commutative graded algebra (CGA) structure that extends the algebra structure on A, respects the A-dgmodule structure, and such that $(sq) \cdot (sq') = 0$, for $q, q' \in Q$. We will call this structure the *semi-trivial CGA structure* on the mapping cone $A \oplus_f sQ$ (see [9, Section 4]). The following result is very useful to detect when this CGA structure is, in fact, a CDGA structure.

Definition 2.4. *Let A be a CDGA. An A-dgmodule morphism $f\colon Q \to A$ is* balanced *if :*

$$f(x)y = xf(y) \quad \textit{for all } x, y \in Q. \tag{4}$$

The importance of this notion comes from the following proposition.

Proposition 2.5. *Let Q be an A-dgmodule and $f\colon Q \to A$ be an A-dgmodule morphism. If f is balanced then the mapping cone $C(f) = A \oplus_f sQ$ endowed with the semi-trivial CGA structure is a CDGA.*

Proof. The only non trivially verified condition for $C(f)$ being a CDGA is the Leibniz rule for the differential, which is a consequence of (4). See the proof of [4, Proposition 2.2] for more details. □

Proposition 2.6. *Let A be a connected CDGA and let $f\colon Q \to A$ be an A-dgmodule morphism. Let p and N be natural integers such that $Q^{<p} = 0$ and $N \leq 2p-3$. Then the semi-trivial CGA structure on the mapping cone $C(f)$ induces a CDGA structure on $\tau^{\leq N}(C(f))$, and*

$$A \to \tau^{\leq N}(C(f))$$

is a CDGA morphism.

Proof. Analogous argument as for the proof of Proposition 2.5. See the proof of [9, lemme 4.5] for more details. □

Remark 2.7. *In the rest of this paper, when a mapping cone is equipped with a CDGA structure it will be understood that it comes from the semi-trivial structure.*

2.5. **Homotopy kernel.** In this section we recall the notion of homotopy kernel and some of its properties.

Definition 2.8. *Let $f\colon M \to N$ be a morphism of A-dgmodules. The* homotopy kernel *of f is the A-dgmodule mapping cone*

$$\text{hoker } f := s^{-1}N \oplus_{s^{-1}f} M,$$

which comes with an obvious map

$$\text{hoker } f \to M \ ; \ (s^{-1}n, m) \mapsto m.$$

The following result is a consequence of the five lemma and justifies the terminology "homotopy kernel".

Proposition 2.9. *Let $f\colon M \to N$ be a surjective morphism of A-dgmodules. Then the morphism*

$$\begin{array}{rccc} \varphi: & \ker f & \xrightarrow{\simeq} & \text{hoker } f \\ & m & \longmapsto & (0,m) \end{array}$$

is an A-dgmodule quasi-isomorphim.

3. Lefschetz duality for manifolds with boundary

The aim of this section is to prove Proposition 3.1 below, which is a first step towards the description of the rational homotopy type of the complement of a subpolyhedron in a manifold with boundary.

Let W be a closed connected oriented triangulated manifold of dimension n with boundary and let $f : K \hookrightarrow W$ be the inclusion of a connected subpolyhedron of dimension k in W. Denote by ∂W the boundary of W and set

$$\partial_W K := K \cap \partial W. \tag{5}$$

In this section we will construct a dgmodule model of $W \setminus K$, extending [11, Theorem 6.3] to manifolds with boundary. Consider the diagram

$$\begin{array}{ccc} W & \xleftarrow{\quad f \quad} & K \\ \uparrow & & \uparrow \\ \partial W & \xleftarrow[\partial f]{} & \partial_W K, \end{array} \tag{6}$$

which after applying the A_{PL} functor gives

$$\begin{array}{ccc} A_{PL}(W) & \xrightarrow{A_{PL}(f)} & A_{PL}(K) \\ \downarrow & & \downarrow \\ A_{PL}(\partial W) & \longrightarrow & A_{PL}(\partial_W K). \end{array} \tag{7}$$

Recall that for a map of spaces $Y \to X$, we set

$$A_{PL}(X, Y) = \ker(A_{PL}(X) \to A_{PL}(Y)).$$

The inclusion of pairs

$$i\colon (K, \partial_W K) \hookrightarrow (W, \partial W)$$

induces an $A_{PL}(W)-$dgmodule morphism

$$A_{PL}(i)\colon A_{PL}(W, \partial W) \to A_{PL}(K, \partial_W K). \tag{8}$$

Using our notation for mapping cones, suspension and linear duals from Section 2.4, consider the map

$$s^{-n}\#A_{PL}(i)\colon s^{-n}\#A_{PL}(K, \partial_W K) \to s^{-n}\#A_{PL}(W, \partial W)$$

and its mapping cone

$$C\left(s^{-n}\#A_{PL}(i)\right) \;=\; s^{-n}\#A_{PL}(W, \partial W) \oplus_{s^{-n}\#A_{PL}(i)} ss^{-n}\#A_{PL}(K, \partial_W K) \tag{9}$$

with the inclusion

$$\iota\colon s^{-n}\#A_{PL}(W, \partial W) \hookrightarrow C\left(s^{-n}\#A_{PL}(i)\right). \tag{10}$$

Since $(W, \partial W)$ is an oriented compact manifold of dimension n, Poincaré duality induces a quasi-isomorphism of $A_{PL}(W)-$dgmodules

$$\Phi_W\colon A_{PL}(W) \xrightarrow{\simeq} s^{-n}\#A_{PL}(W, \partial W) \tag{11}$$

(see (12) in the proof of Proposition 3.1 below for an explicit description of Φ_W.)

Proposition 3.1. *The map*

$$A_{PL}(W) \to A_{PL}(W \setminus K)$$

is weakly equivalent in the category of $A_{PL}(W)$-dgmodules to the map

$$\iota \circ \Phi_W\colon A_{PL}(W) \to C(s^{-n}\#A_{PL}(i))$$

where $C(s^{-n}\#A_{PL}(i))$ is the mapping cone (9)*, ι is from* (10)*, and Φ_W is from* (11)*.*

Proof. First we review from [11, Section 4] a variation of the functor A_{PL} defined on ordered simplicial complex and having an improved excision property. Recall from [6, Chapter 10] that A_{PL} is actually defined first on simplicial sets. Consider the category, $\mathcal{K}$, of ordered simplicial complexes. To any ordered simplicial complex, K, we can associate naturally a simplicial set, $K_\bullet$, whose non-degenerate simplices are exactly the simplices of K (see [5, p.108]). Define the functor

$$\widehat{A_{PL}}\colon \mathcal{K} \to ADGC; K \to A_{PL}(K_\bullet).$$

This functor verifies the two following properties (see [11, Section 4]):

(1) $A_{PL}(|K|) \simeq \widehat{A_{PL}}(K)$ naturally for every ordered simplicial complex (where $|K|$ is the geometric realization).
(2) *Strong excision property* : Let (K, L) be a pair of ordered simplicial complexes. Let $K' \subset K$ a sub-complex and $L' = K' \cap L$. If $K' \cup L = K$ then the inclusion $j\colon (K', L') \hookrightarrow (K, L)$ induces an *isomorphism*

$$\widehat{A_{PL}}(j)\colon \widehat{A_{PL}}(K, L) \xrightarrow{\cong} \widehat{A_{PL}}(K', L').$$

(Note that $A_{PL}(j)$ is a quasi-isomorphism by the classical excision property.)

Consider now the triangulated compact manifold W and its subpolyhedron K. Replace those polyhedra W and K by their second barycentric subdivision. Denote by T the star of K in W, which is a regular neighborhood (see [7, chapters 1 and 2]), hence T is a codimension 0 submanifold with boundary and it retracts by deformation onto K. It is clear that the topological closure $\overline{W \setminus T}$ of $W \setminus T$ is homotopy equivalent to $W \setminus K$. Set

$$\begin{aligned} \partial_+ T &= \partial T \cap \partial W \\ \partial_- T &= \overline{(\partial T \cap (W \setminus \partial W))} = T \cap \overline{W \setminus T} \\ \partial_0 T &= \partial_+ T \cap \partial_- T, \end{aligned}$$

which gives a decomposition of the boundary of T,

$$\partial T = \partial_+ T \cup_{\partial_0 T} \partial_- T.$$

Our next goal is to set up Diagram (13) below. Let us fix an arbitrary order on the vertices of the simplicial complex W such that W and the subpolyhedron T, ∂T, $\partial_+ T$, $\partial_- T$, $\partial_0 T$, K and $\partial_W K$ turn into ordered simplicial complexes. We can apply to them the functor $\widehat{A_{PL}}$ which is naturally quasi-isomorphic to A_{PL}. To prove the result, it suffices to show that the mapping cone $C(s^{-n}\#\ \widehat{A_{PL}}(i))$ is a model of $\widehat{A_{PL}}(W)$-dgmodule of $\widehat{A_{PL}}(\overline{W \setminus T})$. To ease notations, in the rest of this proof we will write A_{PL} instead of $\widehat{A_{PL}}$.

The inclusion of the pair

$$(T, \partial T) \hookrightarrow (W, \overline{W \backslash T} \cup \partial W)$$

induces by the strong excision property above an isomorphism

$$A_{PL}(W, \overline{W \backslash T} \cup \partial W) \xrightarrow{\cong} A_{PL}(T, \partial T).$$

Denote by n the dimension of W. By Poincaré duality of the pair $(W, \partial W)$, there exists an orientation

$$\epsilon_W : A_{PL}(W, \partial W) \to s^{-n}\mathbb{Q},$$

i.e. a morphism of cochain complexes that induces an isomorphism in cohomology in degree n. Using this morphism we can define a morphism of $A_{PL}(W)$-dgmodules

$$\begin{array}{rccc} \Phi_W \colon & A_{PL}(W) & \longrightarrow & s^{-n}\#A_{PL}(W, \partial W) \\ & \alpha & \longmapsto & \left(\Phi_W(\alpha) \colon \beta \mapsto \epsilon_W(\alpha\beta)\right), \end{array} \tag{12}$$

which is a quasi-isomorphism by Poincaré duality of the pair $(W, \partial W)$. The composition

$$\epsilon_T : A_{PL}(T, \partial T) \cong A_{PL}(W, \overline{W \backslash T} \cup \partial W) \xrightarrow{A_{PL}(incl)} A_{PL}(W, \partial W) \xrightarrow{\epsilon_W} s^n\mathbb{Q}$$

induces an isomorphism in cohomology in degree n. Define

$$\begin{array}{rccc} \Phi_T \colon & A_{PL}(T) & \longrightarrow & s^{-n}\#A_{PL}(T, \partial T) \\ & \alpha & \longmapsto & \left(\Phi_T(\alpha) \colon \beta \mapsto \epsilon_T(\alpha\beta)\right) \end{array}$$

which is a quasi-isomorphism of $A_{PL}(W)$-dgmodules by Poincaré duality of the pair $(T, \partial T)$. Also, using the quasi-isomorphism above and the five lemma, it is not difficult to see that the morphism

$$\begin{array}{rccc} \tilde{\Phi}_T \colon & A_{PL}(T, \partial_- T) & \longrightarrow & s^{-n}\#A_{PL}(T, \partial_+ T) \\ & \alpha & \longmapsto & \left(\tilde{\Phi}_T(\alpha) \colon \beta \mapsto \epsilon_T(\alpha\beta)\right) \end{array}$$

is a quasi-isomorphism of $A_{PL}(T)$-dgmodules, hence of $A_{PL}(W)$-dgmodules.

The inclusion

$$(K, \partial_W K) \hookrightarrow (T, \partial_+ T)$$

is a homotopy equivalence and induces a weak equivalence of $A_{PL}(W)$-dgmodules

$$A_{PL}(T, \partial_+ T) \xrightarrow{\simeq} A_{PL}(K, \partial_W K).$$

By the strong excision property, the inclusion


$$(T, \partial_- T) \hookrightarrow (W, \overline{W \backslash T})$$

induces an isomorphism

$$A_{PL}(W, \overline{\overline{W \backslash T}}) \xrightarrow{\cong} A_{PL}(T, \partial_- T).$$

Combining all these morphisms we get the following commutative diagram of $A_{PL}(W)$-dgmodules

(13)



and the two top lines are short exact sequences.

Properties of mapping cones and of short exact sequences imply that, in the category of $A_{PL}(W)$-dgmodules, the morphism

$$A_{PL}(j)\colon A_{PL}(W) \to A_{PL}(\overline{W\backslash T}) \tag{14}$$

on the top right of (13) is equivalent to the map induced between the mapping cones of the horizontal maps of the square $(*)$ in Diagram (13),

$$\mathrm{id}_{A_{PL}(W)} \oplus s0\colon A_{PL}(W) \oplus s0 \to A_{PL}(W) \oplus sA_{PL}(W, \overline{W\backslash T}). \tag{15}$$

Since the vertical maps below the second line of (13) are quasi-isomorphisms, the morphism $\mathrm{id}_{A_{PL}(W)} \oplus s0$ in (15) is equivalent to

$$\iota \circ \Phi_W\colon A_{PL}(W) \to C(s^{-n}\#A_{PL}(i)).$$

The morphism $A_{PL}(j)$ of (14) is clearly equivalent to

$$A_{PL}(W) \to A_{PL}(W\backslash K).$$

This finishes the proof. □

4. Rational model of the complement of a subpolyhedron in a manifold with boundary

In this section we establish the CDGA model of the complement $W \setminus K$ under some unknotting condition, in particular when the codimension of the subpolyhedron is high (Theorem 4.5). We also state a partial CDGA model without unknotting condition (Proposition 4.7.)

Consider the same setting as at the beginning of Section 3, in particular Diagram (6). Suppose given a commutative diagram of CDGAs

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & B \\ {\scriptstyle\alpha}\downarrow & & \downarrow{\scriptstyle\beta} \\ \partial A & \xrightarrow[\partial\varphi]{} & \partial B \end{array} \tag{16}$$

that is a CDGA model of

$$\begin{array}{ccc} W & \xleftarrow{f} & K \\ \uparrow & & \uparrow \\ \partial W & \xleftarrow[\partial f]{} & \partial_W K, \end{array} \tag{17}$$

in other words Diagram (16) is quasi-isomorphic to Diagram (7). Note that in Diagram (16), ∂A and ∂B are just the names of some CDGAs.

The goal of this section is to construct from Diagram (16) a CDGA model of $A_{PL}(W\backslash K)$.

4.1. **Dgmodule model of the complement $W\backslash K$.** Let $\hat{A}$ be a CDGA such that we have the following zig-zag of quasi-isomorphisms

$$A \xleftarrow[\simeq]{\rho} \hat{A} \xrightarrow[\simeq]{\rho'} A_{PL}(W). \tag{18}$$

The morphism ρ' induces a structure of $\hat{A}$-dgmodule on Diagram (7) and the morphism ρ induces a structure of $\hat{A}$-dgmodule on Diagram (16). From Diagram (16) we deduce an $\hat{A}$-dgmodules morphism between the homotopy kernels of α and β (see Section 2.5)

$$\bar{\varphi} : \text{hoker } \alpha \to \text{hoker } \beta. \tag{19}$$

Note also that by Poincaré duality of the pair $(W, \partial W)$, we have a quasi-isomorphism of A-dgmodules

$$\theta_A \colon A \xrightarrow{\simeq} s^{-n}\#\text{hoker } \alpha.$$

Proposition 4.1. *An $\hat{A}$-dgmodule model of*

$$A_{PL}(W) \to A_{PL}(W\backslash K)$$

is given by the composite

$$A \xrightarrow[\theta_A]{\simeq} s^{-n}\#\mathit{hoker}\ \alpha \xhookrightarrow[\iota]{} C(s^{-n}\#\bar{\varphi})$$

where $C(s^{-n}\#\bar{\varphi})$ is the mapping cone of the $\hat{A}$-dgmodules morphism

$$s^{-n}\#\bar{\varphi} \colon s^{-n}\#\mathit{hoker}\ \beta \to s^{-n}\#\mathit{hoker}\ \alpha.$$

Proof. Since (16) is a CDGA model of (7), hoker α is weakly equivalent as an $\hat{A}$-dgmodule to $A_{PL}(W, \partial W)$ and hoker β is weakly equivalent as an $\hat{A}$-dgmodule to $A_{PL}(K, \partial_W K)$. Hence, the result is a direct consequence of Proposition 3.1. □

Remark 4.2. *If the morphisms α and β are surjective then we can work with the genuine kernel instead of the homotopy kernel.*

The major flaw of the dgmodule model of $W \setminus K$ of Proposition 4.1 is that there is no natural CDGA structure on it. The next proposition is a first step to endow this dgmodule model of $W \setminus K$ with the structure of a CDGA.

Proposition 4.3. *Assume given an $\hat{A}$-dgmodule morphism $\varphi^! \colon Q \to A$ weakly equivalent to*

$$s^{-n}\#\bar{\varphi} \colon s^{-n}\#\textit{hoker } \beta \to s^{-n}\#\textit{hoker } \alpha.$$

Then an $\hat{A}$-dgmodule of $A_{PL}(W) \to A_{PL}(W \setminus K)$ is given by

$$A \hookrightarrow C(\varphi^!)$$

where $C(\varphi^!)$ is the mapping cone $A \oplus_{\varphi^!} sQ$.

Proof. This is a direct consequence of Proposition 4.1. □

Remark 4.4. *The existence of such a morphism $\varphi^!$ is guaranteed if we take for Q a cofibrant $\hat{A}$-dgmodule model of $s^{-n}\#$ hoker β.*

This new dgmodule model $C(\varphi^!) = A \oplus_{\varphi^!} sQ$ of $W \setminus K$ has the advantage that A is a CDGA and therefore, under some dimension hypotheses, the semi-trivial CGA structure on the mapping cone described in Section 2.4 makes it into a CDGA. We develop this in the next section.

4.2. **CDGA model of the complement $W \setminus K$.** We work in the setup of diagrams (16)-(17). Remember also the notion of semi-trivial CDGA structure on a mapping cone from Section 2.4 and the notion of CDGA truncation from Section 2.3. Under some codimension and connectedness hypothesis for the inclusion $f \colon K \hookrightarrow W$ we can construct a CDGA model of $W \setminus K$. More precisely, we have:

Theorem 4.5. *Let W be a compact connected oriented triangulated manifold of dimension n with boundary, and let $K \subset W$ be a subpolyhedron of dimension k. Consider Diagram* (17) *and its CDGA model* (16)*. Let r be an integer such that the induced morphisms on homology $H_*(f; \mathbb{Q})$ and $H_*(\partial f; \mathbb{Q})$ are r-connected, that is $H_{\leq r}(W, K; \mathbb{Q}) = 0$ and $H_{\leq r}(\partial W, \partial_W K; \mathbb{Q}) = 0$.*

Suppose given an A-dgmodule Q weakly equivalent to $s^{-n}\#\text{hoker } \beta$ such that $Q^{<n-k} = 0$ and an A-dgmodules morphism

$$\varphi^! \colon Q \to A \tag{20}$$

weakly equivalent to

$$s^{-n}\#\bar{\varphi}\colon s^{-n}\#\text{hoker }\beta \to s^{-n}\#\text{hoker }\alpha.$$

If

$$r \geq 2k - n + 2, \tag{21}$$

then every truncation $\tau^{\leq n-r-1}(C(\varphi^!))$ *of the mapping cone* $C(\varphi^!) = A \oplus_{\varphi^!} sQ$ *equipped with the semi trivial structure is a CDGA, and the morphism*

$$A \longrightarrow \tau^{\leq n-r-1}(C(\varphi^!))$$

is a CDGA model of the inclusion

$$W\backslash K \hookrightarrow W.$$

Moreover it is always possible to construct an A*-dgmodule* Q *and a morphism* $\varphi^!$ *as in* (20).

This generalizes the main result of [9, Theorem 1.2] to manifolds with boundary. A first direct consequence of this theorem is the following corollary on the rational homotopy invariance of the complement under some connectedness-codimension hypotheses.

Corollary 4.6. *Let* W *be a compact triangulated manifold with boundary and* $K \subset W$ *be a subpolyhedron. Assume that* W *and* ∂W *are 1-connected and that the inclusions*

$$K \hookrightarrow W \text{ and } K \cap \partial W \hookrightarrow \partial W$$

are r*-connected with*

$$r \geq 2(\dim K) - \dim W + 2. \tag{22}$$

Then the rational homotopy type of $W\backslash K$ *depends only on the rational homotopy type of the diagram*

$$\begin{array}{ccc} \partial_W K & \xrightarrow{\partial f} & \partial W \\ \downarrow & & \downarrow \\ K & \xrightarrow[f]{} & W. \end{array}$$

The hypotheses (21) (or equivalently (22)) is called *the unknotting condition* and it cannot be removed as shown in [9, Section 9].

Proof of Theorem 4.5. Let $\hat{A}$ be a CDGA such that we have a zig-zag of CDGA quasi-isomorphisms

$$A_{PL}(W) \xleftarrow[\rho']{\simeq} \hat{A} \xrightarrow[\rho]{\simeq} A.$$

Set $N := 2(n-k)-3$. By Proposition 2.6 (with $p = n-k$), $\tau^{\leq N}C(\varphi^!)$ admits the structure of a CDGA induced by the semi-trivial CGA structure on the mapping cone, and the composite

$$A \xrightarrow{\iota} C(\varphi^!) \longrightarrow \tau^{\leq N}C(\varphi^!)$$

is a CDGA morphism.

We now prove that $H^{>N}(W \setminus K) = 0$ where (co)homology of spaces is understood with coefficients in $\mathbb{Q}$. By excision and the connectedness hypotheses on $H(\partial f)$ and $H(f)$,

$$H_{\leq r}(K \cup_{\partial_W K} \partial W\ ,\ K) \ \cong\ H_{\leq r}(\partial W\ ,\ \partial_W K) \ =\ 0$$

and

$$H_{\leq r}(W\ ,\ K) \ =\ 0.$$

Lefschetz duality and the long exact sequence of the triple $(W\ ,\ K \cup_{\partial_W K} \partial W\ ,\ K)$ give

$$H^{\geq n-r}(W \setminus K) \ \cong\ H_{\leq r}(W\ ,\ K \cup_{\partial_W K} \partial W) \ =\ 0.$$

The unknotting hypothesis (21) implies that $N \geq n-r-1$, therefore $H^{>N}(W \setminus K) = 0$.

By Proposition 4.3, $A \to C(\varphi^!)$ is an $\hat{A}$-dgmodule model of $A_{PL}(W) \to A_{PL}(W\setminus K)$. This implies that

$$H^{>N}(C(\varphi^!)) \cong H^{>N}(W\setminus K) = 0,$$

therefore

$$\text{proj} : C(\varphi^!) \longrightarrow \tau^{\leq N}(C(\varphi^!))$$

is a quasi-isomorphism.

Thus the CDGA morphism

$$A \to \tau^{\leq N}\left(C(\varphi^!)\right)$$

is a model of $\hat{A}$-dgmodules of $A_{PL}(W) \to A_{PL}(W\setminus K)$. We will prove that it is actually a CDGA model.

Take a minimal relative Sullivan model (in the sense of [6, Chapter 14])

$$\begin{array}{ccccc} \hat{A} & \xrightarrow[\simeq]{\rho'} & A_{PL}(W) & \longrightarrow & A_{PL}(W\setminus K) \\ & \searrow & & \overset{\lambda'}{\underset{\simeq}{\nearrow}} & \\ & & (\hat{A}\otimes\Lambda V, D). & & \end{array} \tag{23}$$

By Proposition 4.3, $\hat{A} \rightarrowtail \hat{A}\otimes\Lambda V$ is an $\hat{A}$-dgmodule model of $A \to C(\varphi^!)$. Since $(\hat{A}\otimes\Lambda V, D)$ is a cofibrant $\hat{A}$-dgmodule, we can construct a weak equivalence of $\hat{A}$-dgmodules

$$\lambda\colon \hat{A}\otimes\Lambda V \to C(\varphi^!)$$

making commute the following diagram, where the upper part is of CDGA and the lower part is of $\hat{A}$-dgmodules,

(24)
$$\begin{array}{ccccc}
A_{PL}(W) & \longrightarrow & A_{PL}(W\backslash K) & & \\
\rho'\uparrow\simeq & & \lambda'\uparrow\simeq & & \\
\hat{A} & \rightarrowtail & \hat{A}\otimes\Lambda V & & \\
\rho\downarrow\simeq & & \lambda\downarrow\simeq & \searrow^{\bar{\lambda}=\mathrm{proj}\circ\lambda}\ \simeq & \\
A & \longrightarrow & C(\varphi^!) & \xrightarrow[\mathrm{proj}]{\simeq} & \tau^{\leq N}C(\varphi^!).
\end{array}$$

By Lefschetz duality and the hypothesis on the dimension of K

$$H^{<n-k}(W, W\backslash K) \cong H_{>k}(K, \partial_W K) = 0.$$

By minimality of the Sullivan relative model (23), this implies that $V^{<n-k-1} = 0$. Therefore $(\Lambda^{\geq 2}V)^{\leq N} = 0$ and, since $(\tau^{\leq N}C(\varphi^!))^{>N} = 0$, this implies that the composition

$$\bar{\lambda} : (\hat{A}\otimes\Lambda V, D) \xrightarrow{\lambda} C(\varphi^!) \xrightarrow{\mathrm{proj}} \tau^{\leq N}(C(\varphi^!))$$

is a morphism of CDGA. Thus all the solid arrows in Diagram (24) are of CDGAs. This achieves to prove that $A \to \tau^{\leq N}(C(\varphi^!))$ is a CDGA model of $W\backslash K \hookrightarrow W$, as claimed.

It remains to prove the existence of an A-dgmodule Q and a morphism $\varphi^!$. Since $H^{>k}(\mathrm{hoker}\ \beta) \cong H^{>k}(K, \partial_W K) = 0$, we have $H^{<n-k}(s^{-n}\#\ \mathrm{hoker}\ \beta) = 0$. Therefore there exists a cofibrant A-dgmodule model Q of $s^{-n}\#\ \mathrm{hoker}\ \beta$ such that $Q^{<n-k} = 0$. Since, by Poincaré duality, $s^{-n}\#\ \mathrm{hoker}\ \alpha \simeq A$, there exists an A-dgmodule morphism

$$\varphi^! : Q \longrightarrow A$$

weakly equivalent to $s^{-n}\#\bar{\varphi}$. □

Actually even when the unknotting condition (21) of Theorem 4.5 is not satisfied, we still get a partial model of $W\backslash K$. More precisely we get a CDGA model of $W\backslash K$ up to some degree, i.e. a model of the truncation of $A_{PL}(W\backslash K)$. This is the content of the next proposition.

Proposition 4.7. *Consider the same hypotheses as in Theorem 4.5 except that we do not assume the unknotting condition* (21)

Let $l\colon A(W) \to A(W\backslash K)$ *be a CDGA model of* $A_{PL}(W) \to A_{PL}(W\backslash K)$ *such that* $A(W)$ *and* $A(W\backslash K)$ *are connected. Set* $N = 2(n-k)-3$. *Then the CDGA morphism*

$$A \to \tau^{\leq N}C(\varphi^!)$$

is a CDGA model of the composite

$$\pi\circ l\colon A(W) \hookrightarrow A(W\backslash K) \to \tau^{\leq N}A(W\backslash K).$$

Proof of Proposition 4.7. The proof is very similar to that of Theorem 4.5. The details to change are left to the reader. □

Remark 4.8. *We would have preferred in Proposition 4.7 to state that* $A \to \tau^{\leq N}(C(\varphi^!))$ *is a CDGA model of* $A_{PL}(W) \to \tau^{\leq N}(A_{PL}(W\backslash K))$. *But the latter is not well defined because* $A_{PL}(W\backslash K)$ *is not connected and hence we cannot take its truncation. This is the reason for considering instead a model* $l\colon A(W) \to A(W\backslash K)$ *between* connected *CDGAs.*

Note that $N = n-(2k-n+2)-1$ and therefore, under the unknotting condition $r \geq 2k-n+2$, we have that $N \geq n-r-1$. But, Poincaré duality and the r-connectedness implies that $H^{\geq n-r}(W\backslash K) = 0$. Hence Theorem 4.5 is actually a corollary of Proposition 4.7

5. Rational model of the configuration space of two points in a manifold with boundary

In this section we use the results of Section 4 to describe the rational homotopy type of the configuration space of two points in a compact manifold with boundary under 2-connectedness hypotheses. In particular we prove in Corollary 5.5 that the rational homotopy type of $\mathrm{Conf}(W,2)$ depends only on the rational homotopy type of the pair $(W,\partial W)$ when W and ∂W are 2-connected. We also construct in Theorem 5.4 an explicit CDGA model of $\mathrm{Conf}(W,2)$. Moreover in Theorem 5.8 we describe an elegant CDGA model for $\mathrm{Conf}(W,2)$ when the pair $(W,\partial W)$ admits a pretty surjective model in the sense of [4].

Fix a compact connected orientable manifold of dimension n, W, with boundary ∂W. Let

$$\Delta : W \hookrightarrow W \times W \; ; x \mapsto (x,x)$$

be the diagonal embedding. The configuration space of two points in W is the complementary space

$$\mathrm{Conf}(W,2) := (W \times W)\backslash\Delta(W) = \{(x,y) \in W \times W | x \neq y\}.$$

Notice that the diagonal embedding Δ is such that $\Delta(\partial W) \cong \partial W$ and $\Delta^{-1}(\partial W \times \partial W) = \partial W$. In other words, with the notation of (5),

$$\partial_{W\times W}\left(\Delta(W)\right) = \Delta(\partial W) \cong \partial W.$$

Therefore, according to Corollary 4.6, if W and ∂W are connected enough, then the rational homotopy type of $\mathrm{Conf}(W,2) = W\times W\backslash\Delta(W)$ is determined by the square (25) of Proposition 5.1 below. The goal of the next section is to compute a CDGA model of that square.

5.1. CDGA model of the diagonal embedding of the pair $(W,\partial W)$ into $(W\times W, \partial(W\times W))$.

The goal of this section is to prove the following proposition.

Proposition 5.1. *Let* W *be a compact connected orientable manifold with boundary* ∂W. *Suppose given a CDGA surjective model*

$\beta\colon B \twoheadrightarrow \partial B$ of the inclusion $\partial W \hookrightarrow W$. Then a CDGA model of the square

(25)
$$\begin{array}{ccc} W\times W & \xleftarrow{\quad\Delta\quad} & W \\ \uparrow & & \uparrow \\ \partial(W\times W) & \xleftarrow[\partial\Delta]{} & \partial W. \end{array}$$

where Δ is the diagonal map and $\partial\Delta$ is the composition $\partial W \overset{\Delta}{\hookrightarrow} \partial W \times \partial W \hookrightarrow \partial(W\times W)$ is given by the CDGA square

(26)
$$\begin{array}{ccc} B\otimes B & \xrightarrow{\mu} & B \\ {\scriptstyle\alpha}\downarrow & & \downarrow{\scriptstyle\beta} \\ \frac{B\otimes B}{(\ker\beta\otimes\ker\beta)} & \xrightarrow[\tilde\mu]{} & \partial B \end{array}$$

where μ is the multiplication, α is the projection on the quotient, and $\tilde\mu$ the map induced by $\beta\circ\mu$.

The rest of this section is devoted to the proof of this result and for the rest of it we will use the notations introduced in the proposition. First, notice that, since W is a manifold with boundary, $W\times W$ is also a manifold with boundary

$$\partial(W\times W) = W\times\partial W \cup_{\partial W\times\partial W} \partial W\times W.$$

In other words we have a pushout (and homotopy pushout)

(27)
$$\begin{array}{ccc} \partial(W\times W) & \longleftarrow & W\times(\partial W) \\ \uparrow & \text{pushout} & \uparrow \\ (\partial W)\times W & \longleftarrow & \partial W\times\partial W. \end{array}$$

The key argument to prove Proposition 5.1 is that Diagram (25) is the right upper half of the following diagram

(28)
$$\begin{array}{ccccc} W\times W & \xleftarrow{\quad\Delta\quad} & & & W \\ \uparrow & & & & \uparrow \\ \partial(W\times W) & \longleftarrow & W\times(\partial W) & & \uparrow \\ \uparrow & \text{pushout} & \uparrow & & \uparrow \\ (\partial W)\times W & \longleftarrow & \partial W\times\partial W & \longleftarrow & \partial W \end{array}$$

where the maps are the obvious inclusions and diagonals, and the small left lower square in (28) is the homotopy pushout (27).

Lemma 5.2. *The following diagram is a CDGA model of diagram* (28)

(29)
$$\begin{array}{ccccc} B\otimes B & & \xrightarrow{\quad\mu\quad} & & B \\ \downarrow{\scriptstyle\alpha} & & & & \\ P & \longrightarrow & B\otimes\partial B & & \downarrow{\scriptstyle\beta} \\ \downarrow & \textit{pullback} & \downarrow{\scriptstyle\beta\otimes\mathrm{id}} & & \\ \partial B\otimes B & \xrightarrow[\mathrm{id}\otimes\beta]{} & \partial B\otimes\partial B & \xrightarrow{\mu} & \partial B \end{array}$$

where P is the pullback of the small square, α is the morphism given by the universal property, and μ are the multiplication morphisms.

Proof of Lemma 5.2. Using the classical CDGA models for products and diagonal maps on spaces, the fact that A_{PL} turns homotopy pushout of topological spaces into homotopy pullbacks of CDGAs, that a pullback of CDGA surjections is a homotopy pullback, and standard techniques in rational homotopy theory we get that a CDGA model of Diagram (28) is given by the following diagram, where P' denotes the pullback of the left bottom corner of the square,

(30)
$$\begin{array}{ccccc} A_{PL}(W)\otimes A_{PL}(W) & & \xrightarrow{\quad mult\quad} & & A_{PL}(W) \\ \downarrow & & & & \\ P' & \longrightarrow & A_{PL}(W)\otimes A_{PL}(\partial W) & & \downarrow \\ \downarrow & \text{pullback} & \downarrow & & \\ A_{PL}(\partial W)\otimes A_{PL}(W) & \twoheadrightarrow & A_{PL}(\partial W)\otimes A_{PL}(\partial W) & \longrightarrow & A_{PL}(\partial W). \end{array}$$

This diagram is easily seen to be equivalent to Diagram (29).


□

The following lemma computes the small lower left pullback square in Diagram (29).

Lemma 5.3. *We have a pullback in CDGA*

(31)
$$\begin{array}{ccc} \frac{B\otimes B}{(\ker\beta\otimes\ker\beta)} & \xrightarrow{\overline{id_B\otimes\beta}} & B\otimes\partial B \\ \downarrow{\scriptstyle\overline{\beta\otimes id_B}} & & \downarrow{\scriptstyle\beta\otimes id_{\partial B}} \\ \partial B\otimes B & \xrightarrow[id_{\partial B}\otimes\beta]{} & \partial B\otimes\partial B. \end{array}$$

Proof. Consider the following diagram of CDGA's where the internal square is a pullback and α is the map induced by the universal property:



It is straightforward to check that α is surjective and that

$$\ker \alpha = \ker \beta \otimes \ker \beta.$$


Therefore we have an induced isomorphism

$$\overline{\alpha} \colon \frac{B \otimes B}{\ker \beta \otimes \ker \beta} \xrightarrow{\cong} P.$$


□

Proof of Proposition 5.1. Diagram (25) is the upper right part of Diagram (28), therefore, by Lemma 5.2, a CDGA model of (25) is given by the upper right part of (29). Using Lemma 5.3 which computes the pullback P, we deduce that this CDGA model is (26). □

5.2. **A first CDGA model of** $\mathrm{Conf}(W, 2)$**.** Let $\beta \colon B \twoheadrightarrow \partial B$ be a surjective CDGA model of $i \colon \partial W \hookrightarrow W$. Using the results of Section 4, a CDGA model of $\mathrm{Conf}(W, 2) = W \times W \backslash \Delta(W)$ can be obtained from a CDGA model of

(32)


which, by Proposition 5.1 is given by

(33)


Theorem 5.4. *Let W be a compact triangulated manifold with boundary such that W and ∂W are 2-connected. Let $\beta \colon B \twoheadrightarrow \partial B$ be a surjective CDGA model of $\partial W \hookrightarrow W$ and consider the map*

$$\bar{\mu} \colon \ker \beta \otimes \ker \beta \to \ker \beta$$

induced by the multiplication $\mu\colon B\otimes B\to B$. *Suppose given a* $B\otimes B$*-dgmodule morphism*

$$\delta^!\colon D\to B\otimes B$$

weakly equivalent to

$$s^{-2n}\#\bar{\mu}\colon s^{-2n}\#\ker\beta\to s^{-2n}\#(\ker\beta\otimes\ker\beta)$$

and such that $D^{<n}=0$.

Then every truncation $\tau^{\leq 2n-3}C(\delta^!)$ *of the mapping cone of* $\delta^!$ *admits a semi-trivial CDGA structure and*

$$B\otimes B\to\tau^{\leq 2n-3}C(\delta^!)$$

is a CDGA model of $A_{PL}(W\times W)\to A_{PL}(\mathrm{Conf}(W,2))$.

Proof. Since W and ∂W are 2-connected, we have that the morphisms $\Delta\colon W\hookrightarrow W\times W$ and $\partial\Delta\colon\partial W\hookrightarrow\partial(W\times W)$ are 2-connected. So we are under the hypothesis of Theorem 4.5 with $r=2$, and the result is a direct consequence of it. □

We deduce the rational homotopy invariance of $\mathrm{Conf}(W,2)$.

Corollary 5.5. *Let* W *be a compact manifold with boundary. If* W *and* ∂W *are* 2*-connected then the rational homotopy type of* $\mathrm{Conf}(W,2)$ *depends only of the rational homotopy type of the pair* $(W,\partial W)$.

The rational homotopy invariance of Conf $(W,2)$ when W is a closed 2-connected has been established in [8], and [2] gives partial results in the 1-connected case. When W is not simply-connected, [12] shows that there is no rational homotopy invariance.

Remark 5.6. *If we have a CDGA quasi-isomorphism* $B\xrightarrow{\simeq}B'$ *and* $\delta'^!\colon D'\to B'\otimes B'$ *a* $B'\otimes B'$*-dgmodule morphism which is weakly equivalent as a* $B\otimes B-$*dgmodule morphism to* $s^{-2n}\#\bar{\mu}$ *then it follows immediately from Theorem 5.4 that*

$$B'\otimes B'\to\tau^{\leq 2n-3}C(\delta'^!)$$

is also a CDGA model of $A_{PL}(W\times W)\to A_{PL}(\mathrm{Conf}(W,2))$.

5.3. **A CDGA model of** $\mathrm{Conf}(W,2)$ **when** $(W,\partial W)$ **admits a surjective pretty model.** Let W be a compact manifold of dimension n with boundary ∂W such that both W and ∂W are 2-connected. In this section we will construct an elegant CDGA model of $\mathrm{Conf}(W,2)$ when the pair $(W,\partial W)$ admits a *surjective pretty model* in the sense of [4, Definition 3.1]. Let us recall what this means. Suppose given

(i) a connected Poincaré duality CDGA, P, in dimension n ;
(ii) a connected CDGA, Q;
(iii) a CDGA morphism, $\varphi\colon P\to Q$.

Since P is a Poincaré Duality CDGA there exists an isomorphism of P-dgmodules

$$\theta_P\colon P \xrightarrow{\cong} s^{-n}\#P. \tag{34}$$

Consider the composite

$$\varphi^!\colon s^{-n}\#Q \xrightarrow{s^{-n}\#\varphi} s^{-n}\#P \xrightarrow{\theta_P^{-1}} P, \tag{35}$$

which is a morphism of P-dgmodules. Assume that the morphism

$$\varphi\varphi^!\colon s^{-n}\#Q \to Q$$

is balanced (see Definition 2.4) and consider the CDGA morphism

$$\varphi \oplus \mathrm{id}\colon P \oplus_{\varphi^!} ss^{-n}\#Q \to Q \oplus_{\varphi\varphi^!} ss^{-n}\#Q. \tag{36}$$

When (36) is a CDGA model of the inclusion $\partial W \hookrightarrow W$ we say that it is a *pretty model* of the pair $(W, \partial W)$. If moreover φ is surjective (and hence also (36)) we say that is is a surjective pretty model. Then if we consider the differential ideal

$$I = \varphi^!(s^{-n}\#Q) \subset P, \tag{37}$$

[4, Corollary 3.3] states that the CDGA P/I is a CDGA model of W. In [4] we proved that many compact manifolds admit surjective pretty models as for examples even-dimensional disk bundles over closed manifolds, complements of high codimensional polyhedra in a closed manifold, as well as any compact manifold whose boundary retracts rationally on its half-skeleton (see [4, Definition 6.1].)

The objective in this section is to use this model, P/I, of W to construct an elegant model for $\mathrm{Conf}(W, 2)$, analogous to the one constructed in [8] for configuration spaces in closed manifolds.

Since P is a Poincaré duality CDGA, for any homogeneous basis $\{a_i\}_{0\le i\le N}$ of P, there exists a Poincaré dual basis $\{a_i^*\}_{0\le i\le N}$ characterized by $\epsilon(a_i a_j^*) = \delta_{ij}$ where $\epsilon : P^n \to \mathbb{Q}$ is an orientation of P and δ_{ij} is the Kronecker symbol. Let $\Delta \in (P \otimes P)^n$ be the diagonal class of $P \otimes P$ defined as

$$\Delta = \sum_{i=0}^{N} (-1)^{|a_i|} a_i \otimes a_i^*. \tag{38}$$

Denote by

$$\pi\colon P \to P/I$$

the projection. Taking the image of the diagonal Δ by the projection $\pi \otimes \pi\colon P \otimes P \to P/I \otimes P/I$ we get a *truncated diagonal class*

$$\overline{\Delta} = (\pi \otimes \pi)(\Delta) \in (P/I \otimes P/I)^n. \tag{39}$$

Define the map

$$\overline{\Delta}^!\colon s^{-n}P/I \to P/I \otimes P/I \ ;\ s^{-n}x \mapsto \overline{\Delta} \cdot (1 \otimes x). \tag{40}$$

Lemma 5.7. *The map* $\overline{\Delta}^!\colon s^{-n}P/I \to P/I \otimes P/I$ *defined in* (40) *is a* $P/I \otimes P/I$*-dgmodules morphism.*

Proof. In [8, Lemma 5.1] it is shown that for P a connected Poincaré duality CDGA, the morphism $\Delta^!\colon s^{-n}P \to P \otimes P\ ;\ s^{-n}x \mapsto \Delta(1 \otimes x)$ is a $P \otimes P$-dgmodules morphism.

We have the following commutative diagram

$$\begin{array}{ccc} s^{-n}P & \xrightarrow{\Delta^!} & P \otimes P \\ {\scriptstyle s^{-n}\pi}\downarrow & & \downarrow{\scriptstyle \pi\otimes\pi} \\ s^{-n}P/I & \xrightarrow[\overline{\Delta}^!]{} & P/I \otimes P/I. \end{array}$$

Since P/I is a P-dgmodule generated by $1 \in P/I$, this implies that $\overline{\Delta}^!$ is a $P \otimes P$-dgmodules morphism, and the surjectivity of the morphism $\pi : P \to P/I$ implies that $\overline{\Delta}^!$ is a $P/I \otimes P/I$-dgmodules morphism. □

The main result of this section is the following theorem.

Theorem 5.8. *Let* W *be a* 2 *connected compact manifold of dimension* n *whose boundary is* 2*-connected. Suppose that* $(W, \partial W)$ *admits a surjective pretty model of the form* (36)*, and let* $\overline{\Delta}^!$ *be the* $P/I \otimes P/I$*-dgmodules morphism defined in* (40)*. Then the mapping cone*

$$C(\overline{\Delta}^!) = (P/I \otimes P/I) \oplus_{\overline{\Delta}^!} ss^{-n}P/I,$$

equipped with the semi-trivial structure is a CDGA model of $\mathrm{Conf}(W, 2)$*.*

Before proving the theorem, let us fix some notation and prove a lemma. Set

$$B = P \oplus_{\varphi^!} ss^{-n}\#Q \quad \text{and} \quad \partial B = Q \oplus ss^{-n}\#Q.$$

By hypothesis

$$\beta := \varphi \oplus \mathrm{id}\colon B \twoheadrightarrow \partial B \tag{41}$$

is a surjective CDGA model for the inclusion $\partial W \hookrightarrow W$. Also let $B' := P/I$ and notice that the obvious projection $\pi \oplus 0\colon B \xrightarrow{\simeq} B'$ is a quasi-isomorphism of CDGA. According to Theorem 5.4 and Remark 5.6, we only need to show that $\overline{\Delta}^!$ is equivalent to $s^{-2n}\#\bar{\mu}$, which is the content of the following lemma.

Lemma 5.9. *There exists a* $B \otimes B$*-dgmodules commutative square:*

$$\begin{array}{ccc} s^{-n}P/I & \xrightarrow{\overline{\Delta}^!} & P/I \otimes P/I \\ {\scriptstyle \overline{\theta}_P}\downarrow{\scriptstyle \cong} & & {\scriptstyle \cong}\downarrow{\scriptstyle \overline{\theta}_{P\otimes P}} \\ s^{-2n}\#\ker\beta & \xrightarrow[s^{-n}\#\bar{\mu}]{} & s^{-2n}\#\ker\beta \otimes \ker\beta. \end{array} \tag{42}$$

Proof. By Poincaré duality of the CDGA P, we have a P-dgmodules isomorphism $\theta_P : P \xrightarrow{\cong} s^{-n}\#P$. This morphism induces, by construction of the differential ideal $I \subset P$ (see (35) and (37)), a P-dgmodules isomorphism

$$\bar{\theta}_P \colon P/I \xrightarrow{\cong} s^{-n}\# \ker \varphi.$$

The morphism

$$\beta := \varphi \oplus \mathrm{id} \colon B \twoheadrightarrow \partial B$$

is a surjective CDGA model of $\partial W \hookrightarrow W$. We have an obvious isomorphism $\ker \beta \cong \ker \varphi$ as P-dgmodules. So, we have a P-dgmodules isomorphism (that we will also denote $\bar{\theta}_P$)

$$\bar{\theta}_P \colon P/I \xrightarrow{\cong} s^{-n}\# \ker \beta.$$

An easy computation shows that for $(p,u) \in B = P \oplus ss^{-n}\#Q$ and $x \in P/I$,

$$\bar{\theta}_P((p,u)\cdot x) = (p,u)\bar{\theta}(x).$$

Thus $\bar{\theta}_P$ is a morphism of B-dgmodules and, via the multiplication $\mu \colon B \otimes B \to B$, it is a $B \otimes B$-dgmodules morphism. As a direct consequence we have the $B \otimes B$-dgmodules isomorphism

$$\overline{\theta}_{P\otimes P} \colon P/I \otimes P/I \xrightarrow{\cong} s^{-n}\# \ker \beta \otimes s^{-n}\# \ker \beta \cong s^{-2n}\#(\ker \beta \otimes \ker \beta)$$

By Lemma 5.7, the morphism $\overline{\Delta}^!$ is a $P/I \otimes P/I$-dgmodules morphism, and hence it is also a morphism of $B \otimes B$-dgmodules.

Consider the following diagram of $B \otimes B$-dgmodules

$$\begin{array}{ccc} s^{-n}P/I & \xrightarrow{\quad\overline{\Delta}^!\quad} & P/I \otimes P/I \\ {\scriptstyle \overline{\theta}_P}\downarrow{\scriptstyle \cong} & & {\scriptstyle \cong}\downarrow{\scriptstyle \overline{\theta}_P\otimes\bar{\theta}_P} \\ s^{-2n}\# \ker \beta & \xrightarrow[s^{-n}\#\bar{\mu}]{} & s^{-2n}\# \ker \beta \otimes \ker \beta, \end{array}$$

and let us show that it commutes. Since P/I is a $B \otimes B$-dgmodule generated by the element $1 \in P/I$, it suffices to prove that

$$\bar{\theta}_P \otimes \bar{\theta}_P(\overline{\Delta}^!(s^{-n}1)) = s^{-n}\#\bar{\mu}(\bar{\theta}_P(s^{-n}1)).$$

A straightforward computation shows that this is the case. □

Proof of Theorem 5.8. Since W and ∂W are 2-connected, Lemma 5.9, Remark 5.6 and Theorem 5.4 imply that

$$P/I \otimes P/I \to \tau^{\leq 2n-3}C(\overline{\Delta}^!)$$

is a CDGA model of $\mathrm{Conf}(W,2) \hookrightarrow W \times W$. Moreover, we can verify that the morphism $\overline{\Delta}^!$ is balanced, therefore $C(\overline{\Delta}^!)$ is also a CDGA when equipped with the semi-trivial structure. By the 2-connectedness of the manifold W and for degree reasons we have that $C(\overline{\Delta}^!) \xrightarrow{\simeq} \tau^{\leq 2n-3}C(\overline{\Delta}^!)$ is a CDGA quasi-isomorphism. □

5.4. **A CDGA model for** $\mathrm{Conf}(W,2)$ **when** W **is a disk bundle of even rank over a closed manifold.** We apply the model constructed in Section 5.3 to disk bundles.

Let ξ be a vector bundle of even rank, $2k$, for some $k \geq 2$, over some 2-connected closed manifold, M, of dimension m. Then the disk bundle $D\xi$ is a compact manifold of dimension $m+2k$ with boundary the sphere bundle $S\xi$.

Let Q be a Poincaré duality CDGA model of M, let

$$\Delta_Q \in (Q \otimes Q)^m$$

be a diagonal class for Q, and let

$$e \in Q^{2k} \cap \ker(d_Q)$$

be a representative of the Euler class of ξ. Denote by $(\Delta_Q \cdot (e \otimes 1))^!$ the $Q \otimes Q-$dgmodule morphism

$$(\Delta_Q \cdot (e \otimes 1))^! : s^{-(m+2k)}Q \longrightarrow Q \otimes Q, s^{-(m+2k)}q \longmapsto \Delta_Q \cdot (e \otimes q),$$

which is balanced. Consider the mapping cone

$$Q \otimes Q \bigoplus_{(\Delta_Q \cdot (1 \otimes e))^!} ss^{-(m+2k)}Q$$

which is a CDGA.

Theorem 5.10. *With the notation above, assume that the vector bundle ξ is of even rank $2k \geq 4$, and that the base, M, is a 2-connected closed manifold. Then*

$$Q \otimes Q \bigoplus_{(\Delta_Q \cdot (1 \otimes e))^!} ss^{-(m+2k)}Q$$

is a CDGA model of $\mathrm{Conf}(D\xi, 2)$.

Before proving this theorem, let us first deduce the rational homotopy invariance of that configuration space.

Corollary 5.11. *The rational homotopy type of the configuration space of 2 points in a disk bundle of even rank ≥ 4 over a 2-connected closed manifold depends only on the rational homotopy type of the base and on the Euler class.*

Proof of Corollary 5.11. By the main result of [10], the base of the bundle admits a Poincaré duality CDGA model, Q. Let $e \in Q \cap \ker(d_Q)$ be a representative of the Euler class. By Theorem 5.10, a CDGA model of the configuration space, and hence its rational homotopy type since it is simply connected, depends only on those data. □

Proof of Theorem 5.10. Denote by $\bar{z}$ a generator of degree $2k$ and define the CDGA

$$P := \left(\frac{Q \otimes \wedge \bar{z}}{(\bar{z}^2 - e\bar{z})}, D\bar{z} = 0 \right)$$

which is a Poincaré CDGA in dimension $n = m+2k$. Define the CDGA morphism

$$\varphi : P \longrightarrow Q$$

by $\varphi(q_1 + q_2\bar{z}) = q_1 + q_2 \cdot e$, for $q_1, q_2 \in Q$. Then [4, Theorem 4.1] and its proof establishes that the pair $(D\xi, S\xi)$ admits a surjective pretty model associated to φ. We will then use Theorem 5.8 to establish the model of $\mathrm{Conf}(D\xi, 2)$.

Following the notation of [4, proof of Theorem 4.1], one computes that

$$I = \varphi^!(s^{-n}\#Q) = \Phi^!(s^{-2k}Q) = \bar{z} \cdot Q.$$

We need to compute the truncated diagonal class $\overline{\Delta} \in P/I \otimes P/I$. Let $\{q_i\}$ be a homogeneous basis of Q and let $\{q_i^*\}$ be its Poincaré dual basis. Denote by $\omega \in Q^m$ the fundamental class of Q, so that we have

$$q_i \cdot q_j^* = \delta_{ij} \cdot \omega \mod Q^{<m}.$$

Then

$$\{q_i\} \cup \{q_i \cdot \bar{z}\} \tag{43}$$

is a homogeneous basis of P and $-\omega\bar{z}$ is a fundamental class of P. Then the Poincaré dual basis of (43) is given by

$$\{q_i^* \cdot (e - \bar{z})\} \cup \{-q_i^*\}$$

because of the four equations

$$\begin{aligned}
q_i \cdot q_j^*(e - \bar{z}) &= -\delta_{ij}\omega\bar{z} && \mod P^{<n},\\
q_i \cdot (-q_j^*) &= 0 && \mod P^{<n},\\
(q_i\bar{z}) \cdot (q_j^*(e - \bar{z})) &= q_i q_j^*(\bar{z}e - \bar{z}^2) = 0 && \mod P^{<n},\\
(q_i\bar{z}) \cdot (-q_j^*) &= -q_i q_j^* \bar{z} = -\delta_{ij}\omega\bar{z} && \mod P^{<n}.
\end{aligned}$$

Therefore the diagonal class in P is given by

$$\Delta_P = \sum_i (-1)^{|q_i|}(q_i \otimes q_i^*(e - \bar{z}) - q_i\bar{z} \otimes q_i^*) \in P \otimes P,$$

and, since $I = Q\bar{z}$, the truncated diagonal class is

$$\overline{\Delta}_P = \sum_i (-1)^{|q_i|} q_i \otimes q_i^* e \in P/I \otimes P/I.$$

The diagonal class of Q is

$$\Delta_Q = \sum_i (-1)^{|q_i|} q_i \otimes q_i^* \in Q \otimes Q,$$

therefore, using the canonical isomorphism $P/I \cong Q$, we have

$$\overline{\Delta}_P = \Delta_Q \cdot (1 \otimes e).$$

The theorem is then a direct consequence of Theorem 5.8. □

Note that the total space, $E\xi$, of the vector bundle ξ is homeomorphic to the interior of $D\xi$, and therefore $\mathrm{Conf}(E\xi, 2) \simeq \mathrm{Conf}(D\xi, 2)$. In particular, when the bundle is trivial the above gives a model of $\mathrm{Conf}(M \times \mathbb{R}^{2k}, 2)$. Hence we recover partially the result [1, Theorem1].

Interestingly enough we get different models when the bundle is not trivial. Consider for example the quaternionic Hopf line bundle, η, over S^4, of rank 4. In that case we can take $Q = (\mathbb{Q}[x]/(x^2), d_Q = 0)$, with $\deg(x) = 4$, as a model for S^4 and the Euler class is represented by $e = x$. Using the model of Theorem 5.10 one computes easily that the rational cohomology algebra of $\mathrm{Conf}(D\eta, 2)$ is the same as $H^*(S^4 \vee S^4 \vee S^{11}; \mathbb{Q})$, but $\mathrm{Conf}(D\eta, 2)$ is *not* formal because it admits a non trivial Massey product in degree 11.

By contrast, for the trivial bundle of rank 4 over S^4, $\epsilon = S^4 \times \mathbb{R}^4$, one computes that $\mathrm{Conf}(D\epsilon, 2)$ is formal and its rational cohomology algebra is given by

$$H^*(\mathrm{Conf}(S^4 \times \mathbb{R}^4, 2); \mathbb{Q}) \cong \frac{\wedge(x, x', u)}{(x^2, x'^2, ux - ux')}.$$

with $\deg(x) = \deg(x') = 4$ and $\deg(u) = 7$.

Thus the two compact manifolds $D\eta$ and $D\epsilon$ of dimension 8 are homotopy equivalent but their configuration spaces have different Poincaré series. This is because their boundaries, $\partial D\eta = S^7$ and $\partial D\epsilon = S^4 \times S^3$, are not homotopy equivalent.

5.5. **A CDGA model for** $\mathrm{Conf}(W, 2)$ **when** W **is the complement of a subpolyhedron in a closed manifold.** Let M be a 2-connected closed manifold of dimension n. Let $K \subset M$ be a 2-connected subpolyhedron such that $\dim M \geq 2\dim(K) + 3$. In this section we explain how to build a CDGA model of $\mathrm{Conf}(M \backslash K, 2)$.

Let T be a regular neighborhood of K in M, in other words T is a compact codimension 0 submanifold of M that retracts by deformation on K. Then let W be the closure of $M \backslash T$ in M, which is a compact manifold with boundary $\partial W = \partial T$. The interior of W is homeomorphic to $M \backslash K$. Therefore $\mathrm{Conf}(M \backslash K, 2)$ is homotopy equivalent to $\mathrm{Conf}(W, 2)$.

Let us recall how to build a pretty surjective model of $(W, \partial W)$. By [4, Proposition 5.4] one can construct a surjective CDGA model

$$\varphi : P \twoheadrightarrow Q$$

of $K \hookrightarrow M$, where P is a Poincaré duality CDGA and $Q^{\geq n/2-1} = 0$. As explained in [4, end of Section 3], the main result of [9] implies that the pretty model associated to φ,

$$\varphi \oplus \mathrm{id} : P \bigoplus_{\varphi^!} ss^{-m}\#Q \longrightarrow Q \bigoplus_{\varphi\varphi^!} ss^{-n}\#Q$$

is a CDGA model of $(W, \partial W)$. Therefore, by Theorem 5.8, a CDGA model of $\mathrm{Conf}(M\backslash K, 2)$ is given by

$$P/I \otimes P/I \bigoplus_{\overline{\Delta}^!} ss^{-n}(P/I).$$

Let us illustrate this for the configuration space of a punctured manifold. Let M be a closed 2-connected manifold and set $W = M\backslash\{x_0\}$, with $x_0 \in M$. Let P be a Poincaré duality CDGA model of M with fundamental class

$$\omega \in P^n.$$

Pick a homogeneous basis $\{a_i\}_{0\leq i\leq N}$ of P with $a_0 = 1$ and $a_N = \omega$. Let $\{a_i^*\}$ be the Poincaré dual basis. Then the diagonal class is

$$\Delta = 1 \otimes \omega + (-1)^n \omega \otimes 1 + \sum_{i=1}^{N-1} a_i \otimes a_i^*$$

with $1 \leq \deg(a_i) \leq n-1$, for $1 \leq i \leq N-1$. In that case we can take $I = \mathbb{Q} \cdot \omega$ and we get that

$$\overline{P} = P/(\mathbb{Q} \cdot \omega)$$

is a CDGA model of $M\backslash\{x_0\}$ and the truncated diagonal is

$$\overline{\Delta} = \sum_{i=1}^{N-1} a_i \otimes a_i^*.$$

Thus

$$\overline{P} \otimes \overline{P} \bigoplus_{\overline{\Delta}^!} ss^{-n}\overline{P}$$

is a CDGA model of $\mathrm{Conf}(M\backslash\{x_0\}, 2)$.

References

1. F. R. Cohen and L. R. Taylor, *Configuration spaces: applications to Gelfand-Fuks cohomology*, Bull. Amer. Math. Soc. **84** (1978), no. 1, 134–136. MR 0461522 (57 #1507)
2. Hector Cordova Bulens, *Rational model of the configuration space of two points in a simply connected closed manifold*, To appear in Proc. American Math. Soc.
3. Hector Cordova Bulens, Pascal Lambrechts, and Don Stanley, *Models for configuration spaces in manifolds with boundary*, In preparation.
4. ———, *Pretty rational models for poincaré duality pairs*, Available on arXiv.
5. Edward B. Curtis, *Simplicial homotopy theory*, Advances in Math. **6** (1971), 107–209 (1971). MR 0279808 (43 #5529)
6. Yves Félix, Stephen Halperin, and Jean-Claude Thomas, *Rational homotopy theory*, Graduate Texts in Mathematics, vol. 205, Springer-Verlag, New York, 2001. MR MR1802847 (2002d:55014)
7. J. F. P. Hudson, *Piecewise linear topology*, University of Chicago Lecture Notes prepared with the assistance of J. L. Shaneson and J. Lees, W. A. Benjamin, Inc., New York-Amsterdam, 1969. MR MR0248844 (40 #2094)
8. Pascal Lambrechts and Don Stanley, *The rational homotopy type of configuration spaces of two points*, Ann. Inst. Fourier (Grenoble) **54** (2004), no. 4, 1029–1052. MR 2111020 (2005i:55016)

9. ———, *Algebraic models of Poincaré embeddings*, Algebr. Geom. Topol. **5** (2005), 135–182 (electronic). MR MR2135550 (2006g:55012)
10. ———, *Poincaré duality and commutative differential graded algebras*, Ann. Sci. Éc. Norm. Supér. (4) **41** (2008), no. 4, 495–509. MR 2489632 (2009k:55022)
11. ———, *A remarkable DGmodule model for configuration spaces*, Algebr. Geom. Topol. **8** (2008), no. 2, 1191–1222. MR 2443112 (2009g:55011)
12. Riccardo Longoni and Paolo Salvatore, *Configuration spaces are not homotopy invariant*, Topology **44** (2005), no. 2, 375–380. MR MR2114713 (2005k:55024)
13. Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994. MR MR1269324 (95f:18001)

H.C.B. and P.L.: IRMP, Université catholique de Louvain, 2 Chemin du Cyclotron, B-1348 Louvain-la-Neuve, Belgium
E-mail address: hector.cordova@uclouvain.be

E-mail address: pascal.lambrechts@uclouvain.be

D.S.: University of Regina, Department of Mathematics, College West, Regina, CANADA
E-mail address: stanley@math.uregina.ca